



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

## Singular stochastic nonlinear wave equations

### Citation for published version:

Oh, T 2019, *Singular stochastic nonlinear wave equations*..

### Link:

[Link to publication record in Edinburgh Research Explorer](#)

### General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



Sep. 2019, Bonn

①

## Singular stochastic nonlinear wave equations

### Stochastic nonlinear wave equation (SNLW)

with an additive space-time white noise.

$$\begin{aligned} \text{(SNLW)} \quad & \begin{cases} (\partial_t^2 - \Delta) u + u^k = \xi \\ (u, \partial_t u)|_{t=0} = (u_0, u_1) \in \mathcal{N}^s(\mathbb{T}^d) \end{cases} \end{aligned}$$

$$\text{on } \mathbb{R}_+ \times \mathbb{T}^d, \quad \mathbb{T}^d = (\mathbb{R}/2\pi\mathbb{Z})^d$$

$$\mathcal{N}^s(\mathbb{T}^d) = H^s(\mathbb{T}^d) \times H^{s-1}(\mathbb{T}^d)$$

•  $\xi$  = space-time white noise

= Gaussian process indexed by  $(t, x)$  s.t.

$$\mathbb{E}[\xi(t_1, x_1)\xi(t_2, x_2)] = \delta(t_1 - t_2)\delta(x_1 - x_2)$$

• 1to formulation: With  $v = \partial_t u$ ,

$$d \begin{pmatrix} u \\ v \end{pmatrix} + \begin{pmatrix} 0 & -1 \\ -\Delta & 0 \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} + \begin{pmatrix} 0 \\ u^k \end{pmatrix} dt = \begin{pmatrix} 0 \\ dW \end{pmatrix}$$

(2)

Here,  $W = L^2$ -cylindrical Wiener process

$$W(t) = \sum_{n \in \mathbb{Z}^d} \beta_n(t) e_n$$

$$e_m(x) = \frac{1}{(2\pi)^{d/2}} e^{i m \cdot x}$$

$\{\beta_m\}_{m \in \mathbb{Z}^d}$  = family of mutually independent  
complex-valued Brownian motions

" $\Lambda = \mathbb{Z}^d/2$ " conditioned such that  $\beta_{-m} = \overline{\beta_m}$ ,  $m \in \mathbb{Z}^d$   
convention:  $\text{Var}(\beta_n(t)) = t$ .

$$\text{i.e. } \beta_m = \text{Re } \beta_n + i \text{Im } \beta_n$$

↑ indep real B.M.  $/\sqrt{2}$

$$\Lambda = \bigcup_{k=0}^{d-1} (\mathbb{Z}^k \times \mathbb{Z}_+ \times \{0\}^{d-k-1})$$

Then,  $\{\beta_n\}_{n \in \Lambda}$  is independent.



(3)

## Duhamel formulation (= mild formulation)

We say that  $u$  is a soln to SNLW if  $u$  satisfies

$$u(t) = \partial_t S(t) u_0 + S(t) u_1 - \int_0^t S(t-t') u^*(t') dt' + \underbrace{\int_0^t S(t-t') dW(t')}_{\text{stochastic convolution.}}$$

where  $S(t) = \frac{\sin(t|\nabla|)}{|\nabla|}$ ,  $|\nabla| = \sqrt{-\Delta}$

$$\widehat{S(t)f}(m) = \begin{cases} \frac{\sin(tm)}{|m|} \widehat{f}(m), & m \neq 0 \\ t \widehat{f}(m), & m = 0. \end{cases}$$

In the following, we actually study

$$\begin{cases} (\partial_t^2 + 1 - \Delta) u + u^* = \zeta \\ (u, \partial_t u)|_{t=0} = (u_0, u_1) \end{cases}$$

corresponding to the Klein-Gordon eqn but we simply refer to this as SNLW.

Duhamel formula:

(4)

$$u(t) = 2S(t)u_0 + S(t)u_1 - \int_0^t S(t-t')u^*(t')dt' + \int_0^t S(t-t')dW(t')$$

Now,  $S(t) = \frac{\sin(t\langle \nabla \rangle)}{\langle \nabla \rangle}$ ,  $\langle \nabla \rangle = \sqrt{1-\Delta}$   
 $\langle \cdot \rangle = \sqrt{1-|\cdot|^2}$   
 = Japanese bracket.

The only difference appears at the zeroth freq.  
 but it does not affect well-posedness theory

• Stochastic damped NLW (SdNLW)

$$\begin{cases} (\partial_t^2 + \partial_t + 1 - \Delta)u + u^* = \sqrt{2}\xi \\ (u, \partial_t u)|_{t=0} = (u_0, u_1) \end{cases}$$

Ito formulation:

$$\begin{aligned} d \begin{pmatrix} u \\ v \end{pmatrix} + \left\{ \begin{pmatrix} 0 & -1 \\ 1-\Delta & 0 \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} + \begin{pmatrix} 0 \\ u^* \end{pmatrix} \right\} dt \\ = \begin{pmatrix} 0 \\ -v \end{pmatrix} dt + \begin{pmatrix} 0 \\ \sqrt{2}dW \end{pmatrix} \end{aligned}$$



## Duhamel formulation:

(5)

$$u(t) = \partial_t D(t) u_0 + D(t) (u_0 + u_1) \\ - \int_0^t D(t-t') u^h(t') dt' \\ + \sqrt{2} \int_0^t D(t-t') dW(t'),$$

$$\text{where } D(t) = \frac{e^{-\frac{t}{2}} \sin(t \sqrt{\frac{3}{4} - \Delta})}{\sqrt{\frac{3}{4} - \Delta}}.$$

$\Leftarrow$  In terms of local well-posedness, there is no difference from SNLW (either wave or KG.)

### Why care?

①  $\zeta = dW$  has regularity

$$-\frac{d}{2} - \varepsilon \text{ in } x$$

$$-\frac{1}{2} - \varepsilon \text{ in } t, \quad \text{very rough.}$$

expect

$$\Rightarrow \text{stoch. convolution } \mathbb{F} = \int_0^t \frac{\sin((t-t')\langle \nabla \rangle)}{\langle \nabla \rangle} dW(t')$$

has spatial regularity

$$1 - \frac{d}{2} - \varepsilon \quad (< 0 \text{ for } d \geq 2).$$

⇒ analytically challenging.

⑥

③ SdNLW formally preserves the Gibbs measure  $\rho$ .  
( $k \in 2N+1$ )

$$d\rho = Z^{-1} e^{-H(u,v)} du \otimes dv$$

$$= Z^{-1} e^{-\frac{1}{k+1} \int_{\mathbb{T}^d} u^{k+1} dx} e^{-\frac{1}{2} \|u\|_{H^1}^2} du$$

$$\otimes e^{-\frac{1}{2} \|v\|_{L^2}^2} dv$$

$$= Z^{-1} e^{-\frac{1}{k+1} \int_{\mathbb{T}^d} u^{k+1} dx} d\mu_{k+1}(u) \otimes d\mu_0(v),$$

where

$\int_{\mathbb{T}^d} u^{k+1} dx$  - measure

$H(u,v)$  = Hamiltonian (= energy) for NLW/NLKG

$$= \frac{1}{2} \int_{\mathbb{T}^d} |\nabla u|^2 dx + \frac{1}{2} \int_{\mathbb{T}^d} v^2 dx + \frac{1}{k+1} \int_{\mathbb{T}^d} u^{k+1} dx.$$

$\mu_0$  = Gaussian meas with CM space  $H^0(\mathbb{T}^d)$

$$d\mu_0 = Z_0^{-1} e^{-\frac{1}{2} \|u\|_{H^0}^2} du.$$

$$\Leftrightarrow u = \sum_{n \in \mathbb{Z}^d} \frac{g_n(\omega)}{\langle n \rangle^s} e^{im \cdot x}$$

$\{g_n\}$ , indep standard  $\mathbb{C}$ -valued Gaussian r.v.'s  
s.t.  $g_{-m} = \overline{g_m}$ .

SdNLW

NLW in a vectorial form

(7)

$$d \begin{pmatrix} u \\ v \end{pmatrix} + \left\{ \begin{pmatrix} 0 & -1 \\ 1-\Delta & 0 \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} + \begin{pmatrix} 0 \\ u^k \end{pmatrix} \right\} dt$$

$$= \underbrace{\begin{pmatrix} 0 \\ -v \end{pmatrix} dt} + \underbrace{\begin{pmatrix} 0 \\ \sqrt{2} dW \end{pmatrix}}$$

Ornstein-Uhlenbeck process (for  $v$ ) .

• NLW preserves the Gibbs measure  $f$ .

1-d : Friedlander '85

2-d : Oh-Thomann '17.

• OU preserves the white noise  $\mu_0$  for  $v$

and hence preserves the Gibbs measure

$f(du, dv)$ .

•  $\mathcal{L}$  = generator for SdNLW

$$= \mathcal{L}_1 + \mathcal{L}_2$$

$\mathcal{L}_1$  = generator for NLW

$\mathcal{L}_2$  = generator for OU.



$$f \text{ invariant} \Leftrightarrow L^* f = 0$$

(8)

$$\text{i.e. } \int L F(u, v) df(u, v) = 0$$

$$\Leftrightarrow L_1^* f = 0 \text{ AND } L_2^* f = 0.$$

$$\text{b/c } L^* = L_1^* + L_2^*$$

This can be made rigorous by considering the  
"finite-dimensional" approximation:

$$(SdNLW_N) \quad (\partial_t^2 + \partial_t + 1 - \Delta) U^N + P_{\leq N} \left( (P_{\leq N} U^N)^{\sharp} \right) = \sqrt{2} \xi$$

$\Leftrightarrow$  (i) finite dim'l SdNLW on  $\{|m| \leq N\}$  decoupled  
(ii) linear stochastic damped wave dynamics  
 on high freq

i.e. this preserves

$$\begin{aligned} df_N &= Z_N^{-1} e^{-\frac{1}{k+1} \int_{\mathbb{T}} d(P_{\leq N} u)^{k+1} dx} d\mu_1(u) \otimes d\mu_0(v) \\ &= Z_N^{-1} \left( e^{-\frac{1}{k+1} \int (P_{\leq N} u)^{k+1} dx} d\mu_{1,N}(u) \otimes d\mu_{0,N}(v) \right) \\ &\quad \otimes \underbrace{\left( d\mu_{1,N}^{\perp}(u) \otimes d\mu_{0,N}^{\perp}(v) \right)}_{(ii)} \end{aligned}$$

Main Goal:

Prove local well-posedness of  $SdNLW$  along with a good approximation property by  $SdNLW_N$ .

$\Rightarrow$  Then, we can apply Bourgain's invariant measure argument to obtain

- almost sure global well-posedness of  $SdNLW$
- invariance of the Gibbs measure  $\rho$ .

$$\begin{array}{ccc} p_N & \longrightarrow & p \\ | & & \vdots \\ u_N & \longrightarrow & u \end{array} \quad \leftarrow \text{inv.}$$

See Bourgain '94, '96, Burq-Tzvetkov '08.

Idea: Use formal invariance of  $\rho$  as a replacement of a conservation law.

(• parabolic  $\Phi_3^4$ -model: Hairer-Matetski '18.



$\Rightarrow$  shows SdNLW is a stochastic quantization <sup>(10)</sup> equation (SQE) for

$\Phi_d^{k+1}$  - measure for  $u \otimes$  white noise  $\mu_0$  for  $v$ .

• SdNLW is called the canonical stochastic quantization equation. (Ryong, Saito, Shigemoto '85.)

$\Leftarrow$  hyperbolic  $\Phi_d^{k+1}$  - model.

Parabolic counterpart:

Parabolic  $\Phi_d^{k+1}$  - model / stochastic quantization eqn

$$(\partial_t - \Delta) u + u^{k+1} = \xi \quad \text{on } \mathbb{T}^d$$

•  $d=2$ : Da Prato - Debussche '03.

•  $d=3$ : ( $k=3$ )

Hairer '15: regularity structure

Gubinelli - Imkeller - Perkowski '15:  
paracontrolled distributions  
(Catellier - Chouk '18)

Kupiainen '16: renormalization group method.



In the following, we focus on local well-posedness  
 $\Rightarrow$  We only consider the (second) SNLW.

Chap 1 SNLW in 2-d.

Chap 2: quadratic SNLW in 3-d.

Chap 3: cubic SNLW of Hartree-type in 3-d.  
 (with Mamoru Okamoto)

$\Leftarrow$  I did not get to cover this part.

## Chapter 0: Preliminaries

(SNLW)

$$U(t) = \mathcal{U}_t S(t) U_0 + S(t) U_1 - \int_0^t S(t-t') U^*(t') dt' + \Psi$$

$\Psi$  = stochastic convolution

$$= \int_0^t S(t-t') dW(t')$$

$$= \sum_{m \in \mathbb{Z}^d} e_n \int_0^t \underbrace{\frac{\sin((t-t')\langle m \rangle)}{\langle m \rangle}}_{\text{Wiener integral}} d\beta_n(t')$$

• Wiener integral ( $\mathbb{R}$ -valued case)

Given deterministic  $f \in L^2([a, b])$ ,

define  $I(f) = \int_a^b f dB$  by the left endpoint Riemann sum.

$\Rightarrow I(f)$  is a mean 0 Gaussian r.v.

with  $\text{Var}(I(f)) = \|f\|_{L^2([a, b])}^2$ .

i.e.

$I: L^2([a, b]) \rightarrow L^2(\Omega)$  is an isometry  
(onto its image.)

• The following lemma will be useful in studying regularities of various random distributions

•  $\mathcal{H}_k$  = homogeneous Wiener chaoses of order  $k$ .

$$\mathcal{H}_{\leq k} = \bigoplus_{j=1}^k \mathcal{H}_j.$$

( Think of the  $L^2(\Omega)$ -completion of polynomials  
in  $B$ 's of degree  $\leq k$ .

Lemma 1 (Wiener chaos estimate)

Let  $k \in \mathbb{N}$ . Then, we have

$$\|X\|_{L^p(\Omega)} \leq (p-1)^{k/2} \|X\|_{L^2(\Omega)}$$

for any  $p \geq 2$  and  $X \in \mathcal{H}_{\leq k}$ .

( $\Leftarrow$  Nelson's hypercontractivity of the OV semigroup.)

Lemma 2: Let  $\{X_N\}_{N \in \mathbb{N}}$  and  $X$  be spatially homogenous stochastic processes:  $\mathbb{R}_+ \rightarrow \mathcal{D}'(\mathbb{T}^d)$

i.e. for any  $x_0 \in \mathbb{T}^d$ ,

$$\{X(\cdot, t)\}_{t \in \mathbb{R}_+} \text{ and } \{X(x_0 + \cdot, t)\}_{t \in \mathbb{R}_+}$$

have the same law.

- Suppose that  $\exists k \in \mathbb{N}$  s.t.  $X_N(t)$  and  $X(t)$  belong to  $\mathcal{H}_{\leq k}$  for each  $t \in \mathbb{R}_+$ .

Then,



ii)  $t \in \mathbb{R}_+$ . If  $\exists s_0 \in \mathbb{R}$  such that

$$(*) \quad \mathbb{E} [|\hat{X}(t, m)|^2] \lesssim \langle m \rangle^{-d-2s_0}$$

for any  $m \in \mathbb{Z}^d$ , then

$$X(t) \in W^{s, \infty}(\mathbb{T}^d), \quad s < s_0, \text{ a.s.}$$

↑ Bessel potential space

$$\begin{aligned} \|f\|_{W^{s,p}} &= \|\langle \nabla \rangle^s f\|_{L^p} \\ &= \|\mathcal{F}^{-1}(\langle m \rangle^s \hat{f}(m))\|_{L^p}. \end{aligned}$$

Moreover, if  $\exists \theta > 0$  s.t.

$$\mathbb{E} [|\hat{X}_N(t, m) - \hat{X}(t, m)|^2] \lesssim N^{-\theta} \langle m \rangle^{-d-2s_0}$$

for any  $m \in \mathbb{Z}^d$  and  $N \geq 1$ , then

$$X_N \rightarrow X \text{ in } W^{s, \infty}(\mathbb{T}^d), \quad s < s_0, \text{ a.s.}$$

(ii) Given  $h \in \mathbb{R}$ , define the difference operator  $\delta_h$

$$\text{by } \delta_h X(t) = X(t+h) - X(t).$$

Let  $T > 0$  and suppose (i) holds on  $[0, T]$ .

(15)

If  $\exists \sigma \in (0, 1)$  s.t.

$$\mathbb{E} [ |\delta_h \hat{X}(t, m)|^2 ] \lesssim \langle m \rangle^{-d-2s_0+\sigma} |h|^\sigma$$

for any  $m \in \mathbb{Z}^d$ ,  $t \in [0, T]$ ,  $|h| \leq 1$  with  $0 \leq t+h \leq T$ ,

then,

$$X \in C([0, T]; W^{s, \infty}(\mathbb{T}^d))$$

for  $s < s_0 - \frac{\sigma}{2}$ , a.s.

Furthermore, if  $\exists \theta > 0$  s.t.

$$\begin{aligned} \mathbb{E} [ |\delta_h \hat{X}_N(t, m) - \delta_h \hat{X}(t, m)|^2 ] \\ \lesssim N^{-\theta} \langle m \rangle^{-d-2s_0+\sigma} |h|^\sigma, \end{aligned}$$

for any  $m \in \mathbb{Z}^d$ ,  $t \in [0, T]$ ,  $|h| \leq 1$ , and  $N \geq 1$ ,

then  $X_N$  converges to  $X$  in  $C([0, T]; W^{s, \infty}(\mathbb{T}^d))$

for  $s < s_0$ , a.s.

See Mourrat-Weber-Xu, Oh-Okamoto-Tzvetkov

Idea of the proof:

translation invariance

$$\Rightarrow \mathbb{E} [\hat{X}(t, m) \hat{X}(t, m)] = 0$$

for  $m+m \neq 0$ .

(real-valued setting.)

$$\cdot \mathbb{E}[\hat{X}(t, n) \hat{X}(t, m)]$$

$$= \iint \underbrace{\mathbb{E}[X(t, x) X(t, y)]}_{= F(t, x-y)} \underbrace{e_{-m}(x) e_{-m}(y)}_{= e_{-(m+m)}(x) e_m(x-y)} dy dx$$

$$= \hat{F}(t, m) \int_{\mathbb{T}^d} e^{-i(m+m) \cdot x} dx$$

$$= 0 \text{ for } m+m \neq 0.$$

Suppose  $(*)$  holds. Then, for  $p \gg 1$ ,

Suppress  $t$

$$\| \| X \|_{W^{s, \infty}} \|_{L^p(\Omega)} \stackrel{\text{Sobolev}}{\lesssim} \| \| X \|_{W^{s+\varepsilon, p}} \|_{L^p(\Omega)}$$

$$= \| \| \langle \nabla \rangle^{s+\varepsilon} X(x) \|_{L^p(\Omega)} \|_{L^p_x}$$

$$\stackrel{\text{Lem 1}}{\lesssim} p^{k/2} \| \| \sum_{n \in \mathbb{Z}^d} \langle m \rangle^{s+\varepsilon} \hat{X}(m) e_n(x) \|_{L^2(\Omega)} \|_{L^p_x}$$

$$= \left( \sum_{n \in \mathbb{Z}^d} \langle m \rangle^{2(s+\varepsilon)} |\hat{X}(m)|^2 \right)^{1/2}$$

$$\leq C_p \left( \sum_{n \in \mathbb{Z}^d} \langle m \rangle^{2(s+\varepsilon) - d - 2s_0} \right)^{1/2} < \infty$$

$$\text{iff } s+\varepsilon - s_0 < 0 \quad \text{i.e. } s < s_0. \quad \square$$



Lemma 3:

$$\Psi \in C(\mathbb{R}_+; W^{1-\frac{d}{2}-, \infty}(\mathbb{T}^d)), \text{ a.s.}$$

Moreover,

$$\Psi_N = P_{\leq N} \Psi \rightarrow \Psi$$

$$\text{in } C(\mathbb{R}_+; W^{1-\frac{d}{2}-, \infty}(\mathbb{T}^d)), \text{ a.s.}$$

 $\uparrow$  compact-open topologyPf: We only verify  $\otimes$  in Lemma 2.

$$\mathbb{E} [|\hat{\Psi}(t, m)|^2] = \int_0^t \underbrace{(\sin((t-t')\langle m \rangle))^2}_{\langle m \rangle^2} dt'$$

$$\lesssim_t \langle m \rangle^{-2} = \langle m \rangle^{-d-2(1-\frac{d}{2})} = \langle m \rangle^{-d}.$$

• For temporal regularity (ii), use mean value theorem in time  $\square$ • When  $d=1$ ,

$$\Psi \in C(\mathbb{R}_+; W^{\frac{1}{2}-, \infty}(\mathbb{T}^d)), \text{ a.s.}$$

 $\Rightarrow$  LWP of SNLW is trivial.• For  $d \geq 2$ ,  $\Psi(t)$  is NOT a function.  
only a distribution. $\Rightarrow$   $\exists$  issue in making sense of  $(\Psi(t))^k$   
(and hence  $u^k$ .) $\Rightarrow$  Need to renormalize the nonlinearity

Chap 1: SNLW in 2-d

$$u(t) = \partial_t S(t) u_0 + S(t) u_1 - \int_0^t S(t-t') u^{\sharp}(t') dt' + \Psi.$$

Picard 2<sup>nd</sup> iterate:  $\int_0^t S(t-t') \Psi^{\sharp}(t') dt'$

↑ does NOT make sense.

Triviality:  $u_{\varepsilon}$  = soln to SNLW with regularized noise  $\xi_{\varepsilon} = \eta_{\varepsilon} * \xi$ .

Without a proper renormalization,

$$u_{\varepsilon} \rightarrow \text{lin soln (or 0)}$$

Albenorio-Haba-Russo '96  
Oh-Okamoto-Robert '19.

i.e. renormalization is necessary in order to have a non-trivial limit.

Truncated SNLW:

$$(\partial_t^2 + 1 - \Delta) u_N + u_N^{\sharp} = P_{\leq N} \xi.$$

⇒ Truncated convolution  $\Psi_N = P_{\leq N} \Psi$ .

↑ smooth in  $x$ .

(19)

$$\text{Fix } (t, x) \in \mathbb{R}_+ \times \mathbb{T}^2.$$

$\Psi_N(t, x) = \text{mean 0 Gaussian r.v. with}$

$$\text{variance } \sigma_N(t) = \mathbb{E}[\Psi_N^2(t, x)]$$

$$= \sum_{|n| \leq N} \int_0^t \frac{(\sin(t-t') \langle m \rangle)^2}{\langle m \rangle^2} dt'$$

$$\sim t \log N \rightarrow \infty \text{ as } N \rightarrow \infty.$$

Da Prato - Debussche trick ('02-'03)

McKean '95

Bourgain '96

$$\text{Duhamel} \Rightarrow u_N = v_N + \Psi_N$$

↑ postulate  $v_N$  is smoother.

Binomial:

$$u_N^k = \sum_{j=0}^k \binom{k}{j} \Psi_N^j v_N^{k-j}$$

↑ does NOT have a limit (as  $N \rightarrow \infty$ )

Wick renormalization: Replace  $\Psi_N^j$  by

$$:\Psi_N^j(t, x): = H_j(\Psi_N(t, x); \sigma_N(t))$$

↑  
ptwise operation.

↑ time-dependent



$H_j(x; \sigma) = \text{Hermite poly of deg } j.$

defined by the generating function:

$$G(t, x; \sigma) = e^{tx - \frac{1}{2}\sigma t^2} = \sum_{j=0}^{\infty} \frac{t^j}{j!} H_j(x; \sigma)$$

Prop 4:  $T > 0.$

$\{:\Phi_N^j:\}_{N \in \mathbb{N}}$  is Cauchy in  $L^p(\Omega; C_T W_x^{-\varepsilon, \infty})$   
 $p < \infty$

also almost surely in  $C([0, T]; W^{\varepsilon, \infty}(T^2))$

Pf: We only verify  $\otimes$  in Lemma 2.  
 (unif in  $N \geq 1.$ )

Recall: For mean 0 Gaussian r.v.'s  $f$  and  $g$  with variances  $\sigma_f$  and  $\sigma_g$ , we have

$$\begin{aligned} \mathbb{E}[H_j(f; \sigma_f) H_k(g; \sigma_g)] \\ = \delta_{jk} \cdot j! \{ \mathbb{E}[fg] \}^j. \end{aligned}$$

$\Leftarrow$  follows from computing

$$\int_{\Omega} G(t, f; \sigma_f) G(s, g; \sigma_g) dP$$

$\left. \begin{array}{l} \cdot \text{ directly} \\ \cdot \text{ in terms of Hermite poly} \end{array} \right\} \text{compare coeff.}$

$$\mathbb{E}[\widehat{|\Psi_N^j(t, m)|^2}]$$

(21)

$$= \iint \underbrace{\mathbb{E}[\Psi_N^j(t, x) : \Psi_N^j(t, y)]}_{\text{}} e_n(y-x) dx dy$$

$$= j! \left\{ \mathbb{E}[\Psi_N(t, x) \Psi_N(t, y)] \right\}^j$$

$$= \sum_{\substack{m \in \mathbb{Z}^2 \\ |m| \leq N}} \int_0^t \frac{(A \sin((t-t') \langle m \rangle))^2}{\langle m \rangle^2} dt' e_m(x-y)$$

$$\lesssim \sum_{n_1 + \dots + n_j = n} \frac{1}{\langle m_1 \rangle^2 \dots \langle m_j \rangle^2} \lesssim \frac{1}{\langle m \rangle^{2-\varepsilon}}$$

As before, the time difference in (ii) can be established by MVT  $\square$

Define

$$: u_N^k : = \sum_{j=0}^k \binom{k}{j} \underbrace{:\Psi_N^j:}_{\text{}} v_N^{k-j}$$



$$: u^k : = \sum_{j=0}^k \binom{k}{j} : \Psi^j : v^{k-j}$$

for  $u = \Psi + v$  ↙ smoother.

$\Rightarrow$  Renormalized SNLW (rSNLW)

$$\begin{cases} (\partial_t^2 + 1 - \Delta) u + :u^k: = \xi \\ (u, \partial_t u)|_{t=0} = (u_0, u_1) \end{cases}$$

Really means

$$\begin{cases} (\partial_t^2 + 1 - \Delta) v + \sum_{j=0}^k \binom{k}{j} : \Psi^j : v^{k-j} = 0 \\ (v, \partial_t v)|_{t=0} = (u_0, u_1) \end{cases}$$

Theorem 5 : (Gubinelli-Koch-Oh '18)

Let  $k \geq 2$ . Then, rSNLW is locally well-posed in  $N^s(\mathbb{T})$  for

$$(i) \ k \geq 4 : s \geq s_{\text{crit}} = \max \left( \overset{\text{scaling}}{1 - \frac{2}{k-1}}, \overset{\text{conformal}}{\frac{3}{4} - \frac{1}{k-1}}, 0 \right)$$

$$(ii) \ k = 2, 3 : s > s_{\text{crit}}.$$

Idea : Solve the fixed pt problem (\*\*) for  $v$  by

- ① Strichartz estimates (on lin solns)
- ② Prop 4 on  $: \Psi^j :$
- ③ product estimates.



(i) fractional Leibniz rule.

$$(ii) \quad \frac{1}{p} + \frac{1}{q} \leq \frac{1}{r} + \frac{s}{d}$$

$$\Rightarrow \| \langle \nabla \rangle^{-s} (fg) \|_{L^r} \lesssim \| \langle \nabla \rangle^{-s} f \|_{L^p} \| \langle \nabla \rangle^s g \|_{L^q}$$

A typical term

$$\| \langle \nabla \rangle^{-s} (:\Psi^j: v^{k-j}) \|$$

$$\stackrel{(ii)}{\lesssim} \| \langle \nabla \rangle^{-s} : \Psi^j : \| \| \langle \nabla \rangle^s (v^{k-j}) \|$$

$$\stackrel{(ii)}{\lesssim} \| \langle \nabla \rangle^s v \| \| v \|^{k-j-1}$$

$$u = \Psi + v$$

$$\in \Psi + C([0, T]; H^\sigma(\mathbb{T}^2)), \quad \sigma = s \wedge (1-s) \quad \forall s > 0$$

$$\subset C([0, T]; H^{-s}(\mathbb{T}^2)).$$

Back to page 9: LWP of renormalized SdNLW

$\Rightarrow$  a.s. GWP of renormalized SdNLW  
(= hyperbolic  $\Phi_2^{k+1}$ -model)

and invariance of the Gibbs measure

(by applying Bourgain's inv. meas. argument.

• unique ergodicity (cubic): L. Tolomeo '19.

Q: What about (deterministic)  
global well-posedness?

- In the heat case, one simply estimates  $\partial_t \|v\|_{L^p}^p$ .
- In the dispersive setting, we need to rely on the energy but the situation is much more intricate.

We only consider the defocusing cubic case.

$$(\partial_t^2 + 1 - \Delta) v + \underbrace{v^3 + 3v^2\Psi + 3v:\Psi^2: + :\Psi^3:}_{\text{rough perturbation}} = 0$$

rough perturbation

Two difficulties:

①  $v(t) \notin H^1(\mathbb{T}^2)$ .

$\Rightarrow$  can not use the energy

$$H(v, \partial_t v) = \frac{1}{2} \int |\nabla v|^2 dx + \frac{1}{2} \int (\partial_t v)^2 + \frac{1}{4} \int v^4 dx.$$

• We need to smooth  $v$ .

$\Rightarrow$  I-method in the stochastic setting.

② Even if  $v$  were in  $H^1$ ,  $v$  does not satisfy

(deterministic) NLW., i.e.  $H(v)$  is NOT conserved.

• If a noise is a bit smoother

(think of  $\langle \nabla \rangle^2 \Psi \in C_t L_x^2$ ) s.t.  $v \in C_t H_x^1$

then

$$\partial_t H(v) = \int_{\mathbb{T}^2} \partial_t v \left( \underbrace{(\partial_t^2 + 1 - \Delta) v + v^3}_{= -(v + \Psi)^3} \right) dx \quad \leftarrow \text{No need for renormalization}$$

$$\sim v^2 \Psi + v \Psi^2 + \Psi^3$$

$v^3$ -term is gone!!

$$\stackrel{C-S}{\lesssim} (H(v))^{1/2} \left( \|\Psi\|_{C_T L_x^\infty}^2 \int v^4 dx + \|\Psi\|_{C_T L_x^6}^6 \right)^{1/2}$$

$$\leq C(T, \Psi) H(v)$$

$\Rightarrow$  Gronwall. (Burq-Tzvetkov '14 in the random data well-posedness theory on 3-d cubic NLW)



• I-method (= method of almost conservation law) (26)


• CKSTT '02

(after Bougain's high-low method '98)

Let  $N \in \mathbb{N}$ . Define

$0 < s < 1$

$$m_N(m) = \begin{cases} 1, & |m| \leq N \\ \frac{N^{1-s}}{|m|^{1-s}}, & |m| > 2N \end{cases}$$



Set

$I = I_N$  = Fourier multiplier op with  $m_N$ .

low freq : identity

high freq : fractional integration

def & LP theory

$$\|I f\|_{W^{a+\sigma, p}} \lesssim N^\sigma \|f\|_{W^{a, p}}$$

$\forall 0 \leq \sigma \leq 1-s$   
 $\forall 1 < p \leq \infty$

$$\|f\|_{H^s} \lesssim \|I f\|_{H^1} \lesssim N^{1-s} \|f\|_{H^s}$$

$\Rightarrow$   $Iv$  does not satisfy NLW in 2 ways

① b/c of  $I$

② b/c of the terms with  $\Psi$ .

$$\underline{H(Iv)(t) - H(Iv)(0)}$$

$$= \int_0^t \int (\partial_t Iv) \underbrace{(\partial_t^2 + 1 - \Delta) Iv + (Iv)^3}$$

$$= -I(:(\nu + \Psi)^3:)$$

$$= -I(\nu^3 + 3\nu^2\Psi + 3\nu:\Psi^2: + :\Psi^3:)$$

$$= \int_0^t \int_x (\partial_t Iv) (-I\nu^3 + (I\nu)^3)$$

$\nwarrow$   $I$ -method part  
(commutator part)

$$- 3 \int_0^t \int_x (\partial_t Iv) I(\nu^2\Psi)$$

$$- 3 \int_0^t \int_x (\partial_t Iv) I(\nu:\Psi^2:)$$

$$- \int_0^t \int_x (\partial_t Iv) I(:\Psi^3:)$$

} Gronwall part

$$=: A_1 + A_2 + A_3 + A_4$$

## Theorem 6: (GKO-Tolomeo '18)

(28)

The defocusing cubic rSNLW is globally well-posed in  $H^s(\mathbb{T}^2)$ ,  $s > 4/5$ .

### Sketch of the proof:

#### ① Commutator estimates

Lemma 7:  $k \leq 3$

$$\| (If)^k - I(f^k) \|_{L^2} \lesssim N^{-\frac{1-k(1-s)}{2}} \| If \|_{H^1}^k$$

one of the freq. must be high.

Lemma 8: Given  $r > 0$ ,  $0 < \sigma < 1$ ,

$\exists \eta(r) > 0$  and  $p(r) \gg 1$  s.t.

$$\| (If)(Ig) - I(fg) \|_{L^2} \lesssim N^{r - \frac{1-\sigma}{2}} \| f \|_{H^{1-\sigma}} \| g \|_{W^{-\eta, p}}.$$

Lemma 9:  $k = 0, 1, 2$ .  $\forall r, \exists \eta, p$  s.t.

$$\begin{aligned} & \| I(v^k : \Psi^{3-k} :) - (Iv)^k I(: \Psi^{3-k} :) \|_{L^2} \\ & \lesssim N^{-\frac{1-k(1-s)}{2} + r} \| Iv \|_{H^1}^k \| : \Psi^{3-k} : \|_{W^{-\eta, p}}. \end{aligned}$$

( $\Leftarrow$  Lemmas 7 & 8)



Note: We may lower the regularity by using spacetime estimates.

② On the stochastic term:

Lemma 10:  $p < \infty$

$$\left( \mathbb{E} \|I\Phi\|_{L_{T,x}^p}^p \right)^{1/p} \lesssim p^{1/2} T^{1/2 + 1/p} (\log N)^{1/2}.$$

Pf: Separately estimate  $\underbrace{I P_{\leq N} \Phi}_{= P_{\leq N} \Phi}$  and  $I P_{> N} \Phi$ .

$\Rightarrow$  Fubini & Chebyshev,

$$P \left( \|I\Phi\|_{L_{T,x}^p} > \lambda \right) \lesssim \frac{p^{1/2} T^{1/2 + 1/p} (\log N)^{1/2}}{\lambda^p}.$$

□

For Gronwall part, we have

(30)

Lemma 11 : (i)  $k=0, 1$ ,  $\forall 0 < \theta \leq 1-s$

$$\begin{aligned}
 & \left| \int_{\pi^2} (\partial_t I v) (I v)^k I (\Psi^{3-k}) \right| \\
 & \xrightarrow{L_x^2} \xrightarrow{L_x^4} \lesssim N^\theta (1 + H(I v)^{\frac{3}{4}}) \| \Psi^{3-k} \|_{L_T^\infty W_x^{-\theta, 4}} \\
 & \quad \text{BAD} \quad \text{mapping property of } I.
 \end{aligned}$$

(ii) ( $k=2$ )  $\exists c > 0$  s.t.  $\forall 0 < \gamma < 1/8$ ,

$$\begin{aligned}
 & \left| \int_{t_1}^{t_2} \int_{\pi^2} (\partial_t I v) (I v)^2 (I \Psi) dx dt \right| \\
 & \lesssim \left( 1 + (t_2 - t_1) + \int_{t_1}^{t_2} H(v)^{1+c\gamma} \right) \| I \Psi \|_{L_{(t_1, t_2), x}^{\gamma-1}}
 \end{aligned}$$

( $\Leftarrow$  interpolation & Sobolev).

Putting together, we have

(31)

$$H(Iv)(t) - H(Iv)(t_0)$$

$$= \int_{t_0}^t \int_x (\partial_t Iv) \underbrace{(-Iv^3 + (Iv)^3)}_{\text{Lemma 7}}$$

$$- 3 \int_{t_0}^t \int_x (\partial_t Iv) \underbrace{(I(v^2\Psi) - (Iv)^2 I\Psi)}_{\text{Lemma 9}}$$

$$- 3 \int_{t_0}^t \int_x (\partial_t Iv) (Iv)^2 I\Psi$$

Lemma 11(ii)

$$- 3 \int_{t_0}^t \int_x (\partial_t Iv) \underbrace{(I(v:\Psi^2:) - (Iv)I(:\Psi^2:))}_{\text{Lemma 9}}$$

$$- 3 \int_{t_0}^t \int_x (\partial_t Iv) (Iv) I(:\Psi^2:)$$

$$- \int_{t_0}^t \int_x (\partial_t Iv) I(:\Psi^3:)$$

Commutator terms

Gronwall part

Lemma 11(i)



Given  $T > 0$  and  $\theta > 0$ , define

$$A(N) = \frac{\|I\Psi\|_{L_{T,x}^{\log N}}}{\log N}$$

$$\Omega_{M,\gamma,\theta} = \left\{ \max_{k=1,2} \left\| : \Psi^{3-k} : \right\|_{L_T^\infty W_x^{-\gamma(k), p(k)}} \right. \\ \left. + \max_{k=0,1} \left\| : \Psi^{3-k} : \right\|_{L_T^\infty W_x^{-\theta, 4}} \leq M \right\}$$

Commutator part

Brownian part.

Prop 12:  $\exists \gamma = \gamma(s) > 0$ ,  $\alpha = \alpha(s) > 0$   
 $\theta = \theta(s) > 0$ ,

$\beta = \beta(s) > 0$  with  $\beta < \alpha \leq 1 - 3(1-s)$

s.t. if  $\omega \in \Omega_{M,\gamma,\theta}$ ,  $\frac{H(Iv)(t_0)}{|t_0|} \leq N^\beta$   
 $|t_0| \leq T$ ,

then  $\exists \tau = \tau(M, T, s)$

$\delta = \delta(s) > 0$ ,  $N_0 = N_0(T, s)$

s.t.

$$\underline{H(Iv)(t)} \leq N^\alpha$$

provided that  $\frac{|t - t_0|}{A(N)} \leq \tau$

$$\frac{|t|}{A(N)} < T$$

$$A(N) < N^\delta, \quad N \geq N_0$$

(33)

Pf:  $H(t) - H(t_0) \lesssim \int_{t_0}^t N^{-(1-3(1-s))} H^2$

comm part.

$$+ \sum_{k=1}^2 \int_{t_0}^t N^{-\frac{1-k(1-s)}{2} + \gamma} H^{\frac{k+1}{2}} \|:\Psi^{3-k}:\|_{L_T^\infty W_x^{-\gamma(r), p(r)}}$$

$$+ \sum_{k=0}^1 \int_{t_0}^t N^\theta (1 + H^{3/4}) \|:\Psi^{3-k}:\|_{L_T^\infty W_x^{-\theta, 4}}$$

$$+ \left(1 + (t - t_0) + \int_{t_0}^t H^{1+c\eta}\right) \|I\Psi\|_{L_{T,2}^{\eta^{-1}}}$$

choose  $\eta = (\log N)^{-1}$

$F(t) \geq H(t)$  and  $N^\beta \leq F \leq N^\alpha$ .

(i)  $N^{-(1-3(1-s))} H^2$

$\leq N^{-\alpha} F \cdot F \leq F$ ,  $\alpha \leq 1-3(1-s)$

(ii)  $\underline{k=2}$ :  $N^{-\frac{1-2(1-s)}{2} + \gamma} H^{1/2} H$

$\leq N^{-\frac{\alpha}{2}} F^{1/2} F \leq F$

choose  $\gamma \leq \frac{1-s}{2}$

$\underline{k=1}$ :  $N^{-\frac{1-(1-s)}{2} + \gamma} H$

$\leq H \leq F$ .

$$(iii) \quad N^\theta (1 + H^{\frac{3}{4}})$$

(34)

$$\lesssim N^\theta F^{\frac{3}{4}} = \underbrace{N^\theta F^{-\frac{1}{4}}}_{\lesssim 1} \cdot F$$

if  $\theta \leq \frac{\beta}{4}$  i.e. given  $\beta > 0$ ,  
choose  $\theta \ll 1$ .

$$\Rightarrow H(t) \leq H(t_0) + C_{M,T,S} \left( 1 + \int_0^t F^{1+c(\log N)^{-1}} \right) A \log N$$

• Let  $\beta < \tilde{\beta} < \alpha$

$$\text{consider } F(t) = N^{\tilde{\beta} + A(t-t_0)}$$

$$\cdot F(t_0) = N^{\tilde{\beta}} > N^\beta \geq H(t_0)$$

• Suppose that  $\exists t < T$  s.t.  $t - t_0 < \frac{\alpha - \tilde{\beta}}{A}$

$$\text{and } F(t) \leq H(t)$$

(Otherwise, we would have  
 $H(t) \leq F(t) \leq N^{\tilde{\beta} + \alpha - \tilde{\beta}} = N^\alpha, \quad \forall t - t_0 < \frac{\alpha - \tilde{\beta}}{A}$ )

• By continuity of  $H$  and  $F$ ,

$$\exists t_* \text{ with } t_* - t_0 < \frac{\alpha - \tilde{\beta}}{A}$$

$$\text{s.t. } H(t_*) = F(t_*)$$



$$\Rightarrow N^{\tilde{\beta} + \lambda(t_x - t_0)}$$

(35)

$$\leq N^{\tilde{\beta}} + c \left( 1 + N^{\tilde{\beta}} (1 + c(\log N)^{-1}) \right)$$

$$\times \int_{t_0}^{t_x} N^{\lambda(1 + c(\log N)^{-1})(t - t_0)} dt$$

$$\times A \log N$$

$$\leq N^{\tilde{\beta} + \lambda(t_x - t_0)}$$

$$\times \left\{ \frac{1}{N^{\tilde{\beta} - \beta}} N^{\lambda(t_x - t_0)} + c \frac{A \log N}{N^{\tilde{\beta} + \lambda(t_x - t_0)}} \right.$$

$$+ c A (\log N) N^{\tilde{\beta}} c (\log N)^{-1} \times \frac{1}{N^{\lambda(t_x - t_0)}}$$

$$\times \int_{t_0}^{t_x} N^{\lambda(1 + c(\log N)^{-1})(t - t_0)} dt$$

$$= \frac{N^{\lambda(1 + c(\log N)^{-1})(t_x - t_0)} - 1}{\lambda(1 + c(\log N)^{-1}) \log N} \quad \text{drop}$$

$$\Rightarrow 1 \leq \underbrace{\left( \frac{1}{N^{\tilde{\beta}-\beta} N^{\lambda(t_x-t_0)}} + \frac{c A \log N}{N^{\tilde{\beta}} N^{\lambda(t_x-t_0)}} \right)}_{\leq 1/2 \text{ if } A \leq N^{\tilde{\beta}}} \quad (36)$$

$$+ c A \frac{N^{\tilde{\beta}} c (\log N)^{-1} N^{\lambda c (\log N)^{-1} (t_x - t_0)}}{\lambda (1 + c (\log N)^{-1})}$$

$$= c A \frac{e^{c\tilde{\beta} + c\lambda(t_x-t_0)}}{\lambda (1 + c (\log N)^{-1})} \quad \left( N^{c(\log N)^{-1}} = e^c \right)$$

$$\Rightarrow \frac{1}{2} \leq \frac{c A e^{c\alpha}}{\lambda (1 + c (\log N)^{-1})} \quad t_x - t_0 \leq \frac{\alpha - \tilde{\beta}}{\lambda}$$

$$\leq c' \frac{A}{\lambda} e^{c\alpha} \Rightarrow \textcircled{X} \text{ by choosing } \frac{A}{\lambda} = \kappa \ll 1.$$

$$\Rightarrow H(t) \leq F(t) \leq N^\alpha, \quad \forall t - t_0 \leq \frac{\alpha - \tilde{\beta}}{\lambda}$$

$$= \frac{\tau}{A}.$$

□

Prop 13: (almost a.s. GWP): Let  $s > 4/5$ . (37)

Given  $T, \varepsilon > 0$  (unrelated!!),  $\exists \Omega_{T, \varepsilon} \subset \Omega$

s.t. ①  $P(\Omega_{T, \varepsilon}^c) < \varepsilon$

②  $\forall \omega \in \Omega_{T, \varepsilon}$ ,

$\exists!$  soln  $u$  to the defocusing cubic rSNLW on  $[0, T]$  of the form  $u = v + \Psi$

satisfying  $\|v\|_{L_T^\infty H_x^s} \leq C(s, T, \varepsilon)$

Rmk: a.a.s. GWP  $\Rightarrow$  a.s. GWP (ie GWP as an SPDE)  
Borel-Cantelli

Pf: Fix  $N_0 = N_0(s)$

$$N_{k+1} = N_k^\sigma, \quad \sigma > 1$$

s.t.

⊕

$$N_{k+1}^{2(1-s)} N_k^\alpha + N_k^{2\alpha} \ll N_{k+1}^\beta$$

$$\left( \begin{array}{l} \Rightarrow \text{Need } \beta > 2(1-s) \\ 2(1-s) < \beta < \alpha \leq 1-3(1-s) \\ \Rightarrow \underline{s > 4/5} \end{array} \right)$$



• Suppose  $H(I_{N_k} v)(t) \leq N_k^\alpha$

(38)

Then,

$$H(I_{N_{k+1}} v)(t) \lesssim \|I_{N_{k+1}} \vec{v}\|_{H^1}^2 + \|I_{N_{k+1}} v\|_{L^4}^4$$

$$\lesssim N_{k+1}^{2(1-s)} \|\vec{v}\|_{H^s}^2 + \|I_{N_{k+1}} v\|_{H^{1/2}}^4 \quad \vec{v} = (v, \partial_t v)$$

$$(\|f\|_{H^s} \lesssim \|If\|_{H^1}) \quad \leq \|v\|_{H^{1/2}}^4 \lesssim \|I_{N_k} v\|_{H^1}^4$$

$$\lesssim N_{k+1}^{2(1-s)} H(I_{N_k} v) + H^2(I_{N_k} v)$$

$$\lesssim N_{k+1}^{2(1-s)} N_k^\alpha + N_k^{2\alpha} \stackrel{+}{\ll} N_{k+1}^\beta$$

$$\stackrel{++}{\Rightarrow} H(I_{N_{k+1}} v)(t) \leq N_{k+1}^\beta$$

Let  $\Omega_\Lambda(N) = \{A(N) \leq \Lambda\}$

$\Rightarrow$  By Lemma 10,

$$P(\Omega_\Lambda(N)^c) \leq \frac{C^{\log N} T^{\frac{\log N}{2} + 1}}{\Lambda^{\log N}} \lesssim N^{\underbrace{\log C_{T,s} - \log \Lambda}_{< 0}}$$

Define  $\Omega_\Lambda = \bigcap_{k \in \mathbb{Z}_{\geq 0}} \Omega_\Lambda(N_k)$

by choosing  
 $\Lambda \gg 1$

$$\Rightarrow P(\Omega_\Lambda^c) \leq \sum_k N_k^{\log C_{T,s} - \log \Lambda} \lesssim N_0^{\log C_{T,s} - \log \Lambda}$$

$\rightarrow 0$  as  $\Lambda \rightarrow \infty$ .

Choose  $M(T, \varepsilon), L(T, \varepsilon) \gg 1$  s.t.

(39)

$$P(\Omega_L^c) + P(\Omega_{M, \sigma, \theta}^c) < \varepsilon$$

Choose  $k_0 \gg 1$  s.t.

$$\cdot L < N_{k_0}^\beta$$

$$\cdot H(I_{N_{k_0}} v)(0) \leq N_{k_0}^\beta$$

Prop 12

$$H(I_{N_{k_0}} v)(t) \leq N_{k_0}^\alpha, \quad \forall t < \frac{\tau}{A(N_{k_0})}$$

$$\text{With } t = \frac{\tau}{L}$$

(Recall  $A(N_k) \leq L$   
on  $\Omega_L$ )

(++)

$$\Rightarrow H(I_{\underline{N_{k_0+1}}} v)\left(\frac{\tau}{L}\right) \leq N_{k_0+1}^\beta$$

Prop 12

$$\Rightarrow H(I_{N_{k_0+1}} v)\left(2 \frac{\tau}{L}\right) \leq N_{k_0+1}^\alpha$$

(++)

$$\Rightarrow H(I_{\underline{N_{k_0+2}}} v)\left(2 \frac{\tau}{L}\right) \leq N_{k_0+2}^\beta$$

Prop 12

$\Rightarrow \dots$

Iterate the argument  $\left\lfloor \frac{L\tau}{\tau} \right\rfloor + 1$  times.

(Note: increment is indep of  $N$ .)

$\Rightarrow$  Given  $t \in [(j-1)\frac{\tau}{\Lambda}, j\frac{\tau}{\Lambda}]$ , we have (40)

$$\begin{aligned} \|\vec{v}(t)\|_{H^s} &\lesssim H(I_{N_{k_0+j-1}} v)^{1/2}(t) \\ &\leq N_{k_0+j-1}^{\alpha/2} \left( \leq N_{k_0 + [\frac{\Lambda T}{\tau}] - 1}^{\alpha/2} \right) \end{aligned}$$

$k_0, \alpha, \Lambda, \tau$  depend only on  $s, T$ , and  $\varepsilon$ .

□

Rmk: A standard application of the I-method:

Given  $T \gg 1$ , choose  $N = N(T) \gg 1$ .

but in our argument, we needed to choose

an increasing seq  $N_{k_0+j}$ ,  $j = 0, 1, \dots, [\frac{\Lambda T}{\tau}]$

$\Leftarrow$  much more subtle.

J. Følano on BBM '18.

- GWP of the defocusing cubic SNLW on  $\mathbb{R}^2$   
L. Tolomeo '18.



## Chap 2: quadratic SNLW in 3-d

(41)

$$\begin{aligned} (\partial_t^2 + 1 - \Delta) u + u^2 &= \xi \\ (u, \partial_t u)|_{t=0} &= (u_0, u_1) \end{aligned} \quad \text{on } \mathbb{R}_+ \times \mathbb{T}^3.$$

$d=3 \Rightarrow \xi$  has spatial regularity  $-\frac{3}{2} - \varepsilon$ .

• Parabolic case:

(SNLH)  $(\partial_t + 1 - \Delta) u + u^2 = \xi$

Da Prato-Debussche:  $u = v + \eta$

$$\eta = \mathcal{I}(\xi)$$

vertex  $\bullet = \xi$

edge = Duhamel integral operator  $\mathcal{I}$ .  
 $= (\partial_t + 1 - \Delta)^{-1}$

• The heat Duhamel integral operator  $\mathcal{I}$  gains 2 derivatives (with  $dt'$ ) but with  $dt(t')$ , only 1 derivative ( $\Leftarrow$  2 derivatives in the second moment.)

$$\Rightarrow \eta \sim -\frac{1}{2} -$$

We only keep track of spatial regularities.

Rules

- A product of functions of reg.  $S_1$  and  $S_2$  is defined if  $S_1 + S_2 > 0$ . When  $S_1 > 0$  and  $S_1 \geq S_2$ , the resulting prod has regularity  $S_2$ .
- A product of stochastic objects (not depending on the unknown) is always well defined, possibly with a renormalization. The product of stoch. obj. of reg  $S_1$  and  $S_2$  has regularity

$$\min(S_1, S_2, S_1 + S_2)$$

$$\begin{aligned}
 (\text{SNLH}) \stackrel{D-D}{\Rightarrow} (2_t + 1 - \Delta) V &= -(V + 1)^2 \\
 &= -V^2 - 2V1 - \underset{\substack{\uparrow \\ -1-}}{1^0} \\
 &\quad \text{renorm } 1^2 \rightsquigarrow V
 \end{aligned}$$

$$\Rightarrow \text{expect } V \sim 1- = (-1-) + 2.$$

$$\begin{array}{cc}
 V & 1 \\
 1- & -\frac{1}{2}-
 \end{array}
 \quad (1-) + (-\frac{1}{2}-) > 0$$

$\Rightarrow$  The product  $V1$  and hence all the terms on RHS are well defined.

$\Rightarrow$  run a contraction argument in  $C_T C_x^{1-}$

Back to the wave case:

$$\varphi = I(\xi) \sim -\frac{1}{2} -$$

$$\text{Here, } I = (\partial_t^2 + 1 - \Delta)^{-1}$$

(corresp to the forward fund. soln.)

$$\text{Wick power: } \varphi_N = \varphi_N^2 - \sigma_N \text{ with } \varphi_N = P_{\leq N} \varphi$$

$$\sigma_N = \mathbb{E}[\varphi^2(t, x)] \sim t N.$$

$$\Rightarrow \varphi_N \rightarrow \varphi \text{ in } C_T W_x^{-1, \infty} \text{ a.s.}$$

2nd order stoch. obj:

$$\gamma = I(\varphi) = \int_0^t \frac{\sin(t-t') \langle \nabla \rangle}{\langle \nabla \rangle} \varphi(t') dt'$$

A naive "parabolic thinking" yields

$$0- = 2(-\frac{1}{2}-) + 1.$$

$\Leftarrow$  BAD. We need to use the explicit product structure and exhibit multilinear dispersive smoothing



Prop 14:  $Y_N = I(V_N) \rightarrow Y^\circ$

in  $C_T W_x^{\frac{1}{2}-\varepsilon, \infty} \cap C_T^1 W_x^{-1-\varepsilon, \infty}$ , a.s.

i.e.  $Y^\circ \sim \frac{1}{2}- \Leftarrow$  extra  $\frac{1}{2}-$  smoothing

Second order expansion:

"at least as smooth as  $Y^\circ$ "

$$u = 1 - Y^\circ + v.$$

$$\begin{aligned} (\text{SNLW}) \Rightarrow (2_t^2 + 1 - \Delta)v &= -(v + 1 - Y^\circ)^2 + v \\ &= -(v - Y^\circ)^2 - 2v1 + \underbrace{21Y^\circ}_{-\frac{1}{2}-} \end{aligned}$$

$\Rightarrow$  expect  $v \sim \frac{1}{2}-$

$$\Rightarrow v1 : (\frac{1}{2}-) + (-\frac{1}{2}-) < 0$$

The product does not make sense.

So, the second order expansion is NOT enough.

- Littlewood-Paley projection:  $P_j$  onto  $\{|m| \sim 2^j\}$ .

$$f = \sum_{\bar{j}=0}^{\infty} P_{\bar{j}} f.$$

- Paraproduct decomposition (Bony '81).

$$fg = f \otimes g + f \oplus g + f \oslash g$$

$$= \sum_{j < k-2} P_j f P_k g + \sum_{|j-k| \leq 2} P_j f P_k g + \sum_{k < j-2} P_j f P_k g.$$

- $f \otimes g =$  para-product of  $g$  by  $f$ . ( $g =$  high freq)

always well defined as a distribution  
of regularity  $\min(S_2, S_1 + S_2)$

- $f \oplus g =$  resonant product

In general, well defined only if  $S_1 + S_2 > 0$ .

$\Leftarrow$  The resonant prod is the source of difficulty  
in making sense of a nonlinearity.

- Set  $\oplus = \otimes + \oplus$ .

Further decomposition:

Paracontrolled ansatz:  $v = X + Y$

$s_1, s_2, \quad 0 < s_1 < s_2$

$$(SNLW2) \quad (\partial_t^2 + 1 - \Delta) X = -2(X + Y - Y^\circ) \otimes 1$$

$$(\partial_t^2 + 1 - \Delta) Y = -(X + Y - Y^\circ)^2 - 2(X + Y - Y^\circ) \otimes 1$$

Why called a paracontrolled ansatz?

A distribution  $f$  is said to be paracontrolled (by a given reference distribution  $g$ ) if  $\exists f'$  s.t.

$$f = f' \otimes g + h$$

$\nwarrow$  smoother

(SNLW2) says  $\square v = \square X + \square Y$  is paracontrolled.

Expect  $X \sim \frac{1}{2}- = (-\frac{1}{2}-) + 1$ .

For now, ignore  $\otimes$  in the  $Y$ -eqn.

$$(X + Y - Y^\circ) \otimes 1 \sim 0-$$

$$\Rightarrow Y \sim 1-$$

$$\Rightarrow Y \otimes 1 \text{ makes sense as long as } s_2 > \frac{1}{2}.$$



$$\text{Y} = \text{Y} \ominus 1$$

$$0- = \left(\frac{1}{2}-\right) + \left(-\frac{1}{2}-\right) \quad \text{without renormalization}$$

We did not expect any further renorm. since it was not needed for SNLH.

In order to prove this, we need to exploit dispersion at a multilinear level.  
(Otherwise, it looks as if there is a log divergence.)

Prop 15:  $\text{Y}_N = \text{Y}_N \ominus 1_N \rightarrow \text{Y} \text{ in } C_T W_x^{-\varepsilon, \infty} \text{ a.s.}$

•  $X \ominus 1$ ?

$$\left(\frac{1}{2}-\right) + \left(-\frac{1}{2}-\right) < 0 \Rightarrow \text{does not make sense as it is.}$$

Idea: Use the structure of  $X$ , i.e.

$$\begin{aligned} X(t) &= \underbrace{\partial_t S(t) X_0 + S(t) X_1}_{= \vec{S}(t)(X_0, X_1)} - 2 \mathcal{I}((X + Y - Y) \ominus 1) \\ &= \vec{S}(t)(X_0, X_1) \end{aligned}$$

Lem 16:  $s_1 > 0$ .  $(X_0, X_1) \in \mathcal{H}^{s_1}(\mathbb{T}^3)$ .

$$Z_N = (\vec{S}(t)(X_0, X_1)) \ominus I_N$$

↓

$$Z = (\vec{S}(t)(X_0, X_1)) \ominus I$$

← Need to include  $Z$  in an enhanced data set.

in  $C_T H_x^{s_1 - \frac{1}{2} - \varepsilon}$ , a.s.

(A straightforward computation with the Wiener chaos estimate and a direct application of Kolmogorov's continuity criterion.

Note: Set of prob 1 depends on the initial data  $(X_0, X_1)$ .

In the parabolic setting, at this point, one would introduce (smoother) commutators:

$$[I((X+Y-Y') \odot I)] \ominus I$$

$$= [(X+Y-Y') \odot I(I)] \ominus I + \boxed{com_1} \ominus I$$

$$= (X+Y-Y') \odot \underbrace{\{I(I) \ominus I\}}_{\text{explicitly known stock obj. of reg } 0-} + \boxed{com_1} \ominus I + com_2$$

explicitly known  
stock obj. of reg 0-.

BAD



Bad News:  $\text{com}_1$  is NOT smooth in the dispersive setting!!

(49)

Main idea: directly study the following paracontrolled operator (and its res product with  $\mathfrak{I}$ ).

Given  $w \in C(\mathbb{R}_+; H^{s_1}(\mathbb{T}^3))$  with  $0 < s_1 < \frac{1}{2}$ ,

define

$$f_{\odot}(w)(t) = \mathcal{I}(w \odot \mathfrak{I})(t)$$

$$= \sum_{j+k=2} \mathcal{I}(P_j w \cdot P_k \mathfrak{I})$$

$$= \sum_{n \in \mathbb{Z}^3} e_n \sum_{n=n_1+n_2} \int_0^t \frac{\sin((t-t')\langle n \rangle)}{\langle n \rangle} \widehat{w}(t', m_1) \widehat{\mathfrak{I}}(t', m_2) dt'$$

$$|n_1| < |m_2|$$

signifies the paraproduct  $\odot$ .

Goal: Make sense of  $f_{\odot}(w) \odot \mathfrak{I}$

with  $w = X + Y - \Upsilon$ .

• Divide  $f_{\odot}$  into good and bad parts

Fix  $\theta > 0$  small

$f_{\odot}^{(1)}$  = restriction of  $f_{\odot}$  onto  $\{|m_1| \geq |m_2|^\theta\}$

$f_{\odot}^{(2)} = f_{\odot} - f_{\odot}^{(1)}$ .



Namely,

$$J_{\leq}^{(1)}(w)(t) = \sum_{n \in \mathbb{Z}^3} e_n \sum_{n=n_1+n_2} \int_0^t \frac{\sin((t-t')\chi(n))}{\langle n \rangle} \widehat{w}(t', m_1) \widehat{f}(t', m_2) dt'$$

$|n_2|^\theta \lesssim |m_1| \ll |m_2|.$

point:  $|m_1|$  is NOT too small.  $\rightarrow |m_1| \sim |m_2|$

$$\langle n \rangle^{\frac{1}{2}+2\varepsilon} \frac{1}{\langle n \rangle} \lesssim \langle m_1 \rangle^{\frac{4\varepsilon}{\theta}} \langle m_2 \rangle^{-\frac{1}{2}-2\varepsilon}$$

$$\lesssim \langle m_1 \rangle^{s_1-\varepsilon} \langle m_2 \rangle^{-\frac{1}{2}-2\varepsilon}$$

by choosing  $\varepsilon = \varepsilon(s_1, \theta) > 0$  suff. small.

Lem 17:  $0 < s_1 < 1/2$ . Given small  $\theta > 0$ ,  
 $\exists$  small  $\varepsilon = \varepsilon(s_1, \theta) > 0$  s.t. given any  
 $\square \in C(\mathbb{R}_+; W^{\frac{1}{2}-\varepsilon, \infty}(\mathbb{T}^3))$ ,

the paracontrolled operator

$$J_{\leq}^{(1), \square} = J_{\leq}^{(1)} \text{ with } f \text{ replaced by } \square$$

belongs to  $L_2 = \mathcal{L}(C_T H_x^{s_1}; C_T H_x^{\frac{1}{2}+2\varepsilon})$

$\uparrow$

following the notation from GKO'18.

(51)

- When  $|m_1| \ll |m_2|^\theta$ , the positive reg of  $w$  does not help.

We directly study  $J_{\odot, \ominus}$ .

$$J_{\odot, \ominus}(w)(t) = J_{\odot}^{(2)}(w) \ominus I(t)$$

$$= \sum_{n \in \mathbb{Z}^3} e_n \int_0^t \sum_{n_1 \in \mathbb{Z}^3} \widehat{w}(t', m_1) A_{n, m_1}(t, t') dt'$$

$$A_{n, m_1}(t, t') = \mathbb{1}_{[0, t]}(t') \sum_{\substack{n - n_1 = n_2 + n_3 \\ |n_1| \ll |m_2|^\theta}} \frac{\sin((t-t')\langle m_1 + m_2 \rangle)}{\langle m_1 + m_2 \rangle} \widehat{f}(t', m_2) \widehat{f}(t, n_3)$$

$(|m_1 + m_2| \sim |n_3|)$  — resonant product

$$= \mathbb{1}_{[0, t]}(t') \sum \frac{\sin((t-t')\langle m_1 + m_2 \rangle)}{\langle m_1 + m_2 \rangle} (\widehat{f}(t', m_2) \widehat{f}(t, n_3) - \mathbb{1}_{m_2 + n_3 = 0} \sigma_{n_2}(t, t'))$$

$$+ \mathbb{1}_{[0, t]}(t') \cdot \mathbb{1}_{m=n_1} \sum_{\substack{n_2 \in \mathbb{Z}^3 \\ |n| \ll |m_2|^\theta}} \frac{\sin((t-t')\langle m + m_2 \rangle)}{\langle m + m_2 \rangle} \sigma_2(t, t')$$

$$=: A_{m, m_1}^{(1)}(t, t') + A_{n, n_2}^{(2)}(t, t')$$

deterministic counter term.

More difficult!!

Here,  $0 \leq t_2 \leq t_1$ ,

$$\sigma_m(t_1, t_2) = \mathbb{E}[\widehat{f}(t_1, m) \widehat{f}(t_2, m)]$$

$$= \frac{\cos((t_1 - t_2)\langle m \rangle)}{2\langle m \rangle^2} t_2 + O\left(\frac{1}{\langle m \rangle^3}\right).$$



(52)

By exploiting dispersion (stationary phase)  
and symmetrization ( $n_2 \leftrightarrow -n_2$ )

i.e. the order of summation matters  
 $\Rightarrow$  only conditionally convergent.

Prop 18:  $S_2 < 1$ .  $\exists$  small  $\theta(S_2) > 0$  and  $\varepsilon > 0$  s.t.

$f_{\otimes, \ominus}$  belongs to the class

$$\mathcal{L}_1 = \mathcal{L} \left( C_T L_x^2 \cap C_T^1 H_x^{+\varepsilon} ; C_T H_x^{S_2-1} \right)$$

Moreover,  $f_{\otimes, \ominus}^N$  (replacing 1 by  $\mathbb{I}_N$ )

converges a.s. to  $f_{\otimes, \ominus}$  in  $\mathcal{L}_1$ .

Final form:

$$\begin{aligned} (\partial_t^2 + 1 - \Delta) X &= -2(X + Y - Y') \otimes \mathbb{I} \\ (\partial_t^2 + 1 - \Delta) Y &= -(X + Y - Y')^2 - 2(X + Y - Y') \otimes \mathbb{I} \end{aligned}$$

(SNLW/3)

$$-2Y \otimes \mathbb{I} + 2 \underline{Y \cdot Y} - 2 \underline{\Delta}$$

$$+ 4 f_{\otimes}^{(1)}(X + Y - Y') \otimes \mathbb{I}$$

$$+ 4 \underline{f_{\otimes, \ominus}(X + Y - Y')}$$

$$(X, \partial_t X, Y, \partial_t Y) \Big|_{t=0} = (X_0, X_1, Y_0, Y_1).$$



Thm 19:  $\frac{1}{4} < s_1 < \frac{1}{2} < s_2 \leq s_1 + \frac{1}{4}$ .

$\exists \theta = \theta(s_2) > 0$  and  $\varepsilon = \varepsilon(s_1, s_2, \theta) > 0$  s.t.

if (i)  $I, Y, \tilde{Y}$  in  $C(\mathbb{R}_+; W_x^{s, \infty})$

$s = -\frac{1}{2} -, \frac{1}{2} -, 0 -$ .

and

$Y \in C'(\mathbb{R}_+; W_x^{-1-\varepsilon})$

(ii)  $Z \in C(\mathbb{R}_+; H_x^{s_1 - \frac{1}{2} - \varepsilon})$ .

(iii)  $f_{\otimes, \oplus} \in \mathcal{L}_1$ ,

NOT random

then the system (SNLW3) is locally well-posed.

More precisely, given  $(X_0, X_1, Y_0, Y_1) \in \mathcal{H}^{s_1} \times \mathcal{H}^{s_2}$ ,

$\exists T$  and unique soln  $(X, Y)$  in the class

$$\Sigma_T^{s_1, s_2} = X_T^{s_1} \times Y_T^{s_2}$$

$$\subset C([0, T]; H^{s_1} \times H^{s_2}) \cap C'([0, T]; H^{s_1-1} \times H^{s_2-2})$$

depending continuously on the enhanced data set.

$$\square = (X_0, X_1, Y_0, Y_1, I, Y, \tilde{Y}, Z, f_{\otimes, \oplus})$$

in the class

$$\begin{aligned} \mathcal{X}_T^{s_1, s_2, \varepsilon} = & \mathcal{H}^{s_1} \times \mathcal{H}^{s_2} \times C_T W_x^{-\frac{1}{2}-\varepsilon, \infty} \times \left( C_T W_x^{\frac{1}{2}-\varepsilon} \cap C_T' W_x^{1-\varepsilon, \infty} \right) \\ & \times C_T W_x^{-\varepsilon, \infty} \times C_T H^{s_1 - \frac{1}{2} - \varepsilon} \times \mathcal{L}_1 \end{aligned}$$

$$u_N = p_N - \dot{Y}_N + \underbrace{X_N + Y_N}$$

$$\downarrow \quad \downarrow \quad \downarrow$$

$$u = p - \dot{Y} + X + Y \text{ in } C_T H_x^{\frac{1}{2}-\varepsilon}$$

Pf of Thm 19: Duhamel formulation.

$$X(t) = \Phi_1(X, Y)(t)$$

$$= \vec{S}(t)(X_0, X_1) - 2 \int_0^t \vec{S}(t-t') [(X+Y-\dot{Y}) \odot 1](t') dt'$$

$$Y(t) = \Phi_2(X, Y)(t)$$

$$= \vec{S}(t)(Y_0, Y_1) - \int_0^t \vec{S}(t-t') [\dots](t') dt'$$

Strichartz estimates:  $0 \leq s \leq 1$ .

We say a pair  $(q, r)$  is  $s$ -admissible

$(\tilde{q}, \tilde{r})$  is dual  $s$ -admis.

(i.e.  $(\tilde{q}, \tilde{r})$  is  $(1-s)$ -admis.)

if  $1 \leq \tilde{q} < 2 \leq q \leq \infty$ ,  $1 < \tilde{r} \leq 2 \leq r < \infty$ ,

$$\frac{1}{q} + \frac{d}{r} = \frac{d}{2} - s = \frac{1}{\tilde{q}} + \frac{d}{\tilde{r}} - 2 \quad \leftarrow \text{scaling condition}$$

$$\frac{1}{q} + \frac{d-1}{2r} \leq \frac{d-1}{4}$$

$$\frac{2}{q} + \frac{d-1}{2\tilde{r}} \geq \frac{d+3}{4}$$

} admissibility cond.

Then, a soln  $u$  to

$$\begin{cases} (\partial_t^2 + 1 - \Delta) u = f \\ (u, \partial_t u)|_{t=0} = (u_0, u_1) \end{cases}$$

satisfies

$$\|(u, \partial_t u)\|_{C_T H_x^s} + \|u\|_{L_T^q L_x^r}$$

$$\lesssim \|(u_0, u_1)\|_{H^s} + \|f\|_{L_T^{\tilde{q}} L_x^{\tilde{r}} + L_x^1 H_x^{s-1}}$$

for  $0 \leq T \leq 1$ .

$(q, r)$ ,  $s$ -admis

$(\tilde{q}, \tilde{r})$ , dual  $s$ -admis.

( $\Leftarrow$  follows from the Strichartz estimates on  $\mathbb{R}^d$  and finite speed of propagation.

Back to the proof of Thm 19: Define the Strichartz space

$$X_T^{S_1} = C_T H_x^{S_1} \cap C_T' H_x^{S_1-1} \cap L_T^p W_x^{S_1-\frac{1}{4}, \frac{p}{3}} \quad (p, \frac{p}{3}), \frac{1}{4}\text{-admis}$$

$$Y_T^{S_2} = C_T H_x^{S_2} \cap C_T' H_x^{S_2-1} \cap L_T^4 W_x^{S_2-\frac{1}{2}, 4} \quad (4, 4), \frac{1}{2}\text{-admis}$$

$$Z_T^{S_1, S_2} = X_T^{S_1} \times Y_T^{S_2}$$



(56)

$$\begin{aligned}
\| \Phi_1(X, Y) \|_{X_T^{s_1}} &\lesssim \| (X_0, X_1) \|_{X^{s_1}} + \| (X+Y-\tilde{Y}) \otimes 1 \|_{L_T' H_x^{s_1-1}} \\
&\lesssim T \| X+Y-\tilde{Y} \|_{L_T^\infty L_x^2} \| 1 \|_{L_T^\infty W_x^{\frac{1}{2}-\varepsilon, \infty}} \\
&\lesssim \| (X_0, X_1) \|_{X^{s_1}} + T (1 + \| (X, Y) \|_{\Sigma_T^{s_1, s_2}}) \\
&\text{if } s_1 - 1 < -\frac{1}{2} - \varepsilon.
\end{aligned}$$

Y-equation:

$$\begin{aligned}
&-2(X+Y-\tilde{Y}) \otimes 1 - 2Y \otimes 1 + 2\tilde{Y} + Z \\
&+ g_{\otimes, \otimes}^{(1)}(X+Y-\tilde{Y}) \otimes 1
\end{aligned}$$

$$\Rightarrow \| \dots \|_{Y_T^{s_1}} \lesssim \| (Y_0, Y_1) \|_{Y^{s_2}} + T (1 + \| (X, Y) \|_{\Sigma_T^{s_1, s_2}})$$

$$\text{if } \frac{1}{2} < s_2 < \min(1, s_1 + \frac{1}{2})$$

Also

$$\begin{aligned}
&\| \int_0^t s(t-t') g_{\otimes, \otimes}(X+Y-\tilde{Y})(t') dt' \|_{Y_T^{s_2}} \\
&\lesssim \| g_{\otimes, \otimes}(X+Y-\tilde{Y}) \|_{L_T' H_x^{s_2-1}} \\
&\lesssim T \| X+Y-\tilde{Y} \|_{L_T^\infty L_x^2 \cap C_T' H_x^{1-\varepsilon}} \\
&\lesssim T (1 + \| (X, Y) \|_{\Sigma_T^{s_1, s_2}}).
\end{aligned}$$

$$\left\| \int_0^t S(t-t') (X+Y-Y')^2(t') dt' \right\|_{Y_T^{S_2}}$$

Strichartz

$$\lesssim \left\| \langle \nabla \rangle^{S_2 - \frac{1}{2}} (X+Y-Y')^2 \right\|_{L_{T,x}^{4/3}}$$

frac. Leib.

$$\lesssim T^{\frac{1}{4}} \left( \left\| \langle \nabla \rangle^{S_2 - \frac{1}{2}} X \right\|_{L_T^{\frac{2}{3}} L_x^{\frac{2}{3}}}^2 + \left\| \langle \nabla \rangle^{S_2 - \frac{1}{2}} Y \right\|_{L_{T,x}^4}^2 + \left\| \langle \nabla \rangle^{S_2 - \frac{1}{2}} Y' \right\|_{L_{T,x}^\infty}^2 \right)$$

Need  
 $\leq S_1 - \frac{1}{4}$

$$\lesssim T^{\frac{1}{4}} \left( 1 + \|(X, Y)\|_{\Sigma_T^{S_1, S_2}}^2 \right)$$

$$\text{if } S_2 \leq \min(1 - \varepsilon, S_1 + \frac{1}{4}).$$

$$\Rightarrow \|\Phi(X, Y)\|_{\Sigma_T^{S_1, S_2}} \lesssim \|(X_0, X_1, Y_0, Y_1)\|_{N^{S_1} \times N^{S_2}} + T^\theta \left( 1 + \|(X, Y)\|_{\Sigma_T^{S_1, S_2}}^2 \right)$$

for some  $\theta > 0$ .

Similarly,

$$\begin{aligned} & \|\Phi(X, Y) - \Phi(\tilde{X}, \tilde{Y})\|_{\Sigma_T^{S_1, S_2}} \\ & \lesssim T^\theta (1 + \|(X, Y)\| + \|(\tilde{X}, \tilde{Y})\|) \|(X, Y) - (\tilde{X}, \tilde{Y})\|_{\Sigma_T^{S_1, S_2}} \end{aligned}$$

$\Rightarrow$  By taking  $T > 0$  suff. small,

$\Phi$  is a contraction on a ball  $B_R \subset \Sigma_T^{S_1, S_2}$

□

# Key point of the proof of Prop 14 on $\hat{Y}$

In using Lemma 2, we bound

$$\mathbb{E}[|\hat{Y}(t, m)|^2]$$

$$= 4 \sum_{\substack{m = n_1 + n_2 \\ n_1 \neq \pm n_2}} \int_0^t \frac{\sin(t-t_1)m}{\langle m \rangle} \int_0^{t_1} \frac{\sin(t-t_2)m}{\langle m \rangle} \Gamma_{m_1}(t_1, t_2) \\ \times \Gamma_{n_2}(t_1, t_2) dt_2 dt_1$$

$$+ \dots_{n_1 = n_2}$$

$$\Gamma_n(t_1, t_2) = \frac{\cos(t_1 - t_2)\langle m \rangle}{2\langle m \rangle^2} t_2 + O\left(\frac{1}{\langle m \rangle^3}\right)$$

$\Rightarrow$  expand sines and cosines in complex exponentials

$$\sum_{\substack{m = n_1 + n_2 \\ n_1 \neq \pm n_2}} \frac{e^{i(\varepsilon_1 + \varepsilon_2)t\langle m \rangle}}{\langle m \rangle^2 \langle m_1 \rangle^2 \langle m_2 \rangle^2} \quad \varepsilon_j \in \{\pm 1\}$$

$$\times \int_0^t e^{-it_1 K_1(\vec{m})} \int_0^{t_1} \frac{1}{t_2^2} e^{-it_2 K_2(\vec{m})} dt_2 dt_1$$

$$K_1(\vec{m}) = \varepsilon_1 \langle m \rangle - \varepsilon_3 \langle m_1 \rangle - \varepsilon_4 \langle m_2 \rangle$$

$$K_2(\vec{m}) = \varepsilon_2 \langle m \rangle + \varepsilon_3 \langle m_1 \rangle + \varepsilon_4 \langle m_2 \rangle$$

integrate in  $t_1$  first.

$$\left| \int_0^t \frac{1}{t_2^2} e^{-it_2 K_2(\vec{m})} \frac{e^{-it K_1(\vec{m})} - e^{-it_2 K_1(\vec{m})}}{-i K_1(\vec{m})} dt_2 \right|$$

$$\leq \frac{C(T)}{(1 + K_1(\vec{m}))}$$




Need to bound

$$I = \sum_{n=n_1+n_2} \frac{1}{\langle m \rangle^2 \langle m_1 \rangle^2 \langle m_2 \rangle^2 (1 + |K_1(\bar{m})|)} \quad \left( \begin{array}{l} \text{WTS} \\ \lesssim \langle m \rangle^{-4+} \end{array} \right)$$

BAD case  $(\varepsilon_1, \varepsilon_3, \varepsilon_4) = (\pm 1, \pm 1, \mp 1)$ . 
 $\left\{ \begin{array}{l} \text{Assume} \\ |m| \gg 1 \\ |n_1| \geq |n_2| \end{array} \right.$

$$|K_1(\bar{m})| = \langle m \rangle + \langle m_2 \rangle - \langle m_1 \rangle$$

Note:  $\langle m_1 \rangle \sim \langle m \rangle + \langle m_2 \rangle$



law of cosines:  $|m|^2 + |m_2|^2 - |m_1|^2 = 2|m||m_2| \cos(\angle(m, m_2))$ .

$$\Rightarrow |K_1(\bar{m})| = \frac{(\langle m \rangle + \langle m_2 \rangle)^2 - \langle m_1 \rangle^2}{\langle m \rangle + \langle m_2 \rangle + \langle m_1 \rangle}$$

$$= \frac{2\langle m \rangle \langle m_2 \rangle + |m|^2 + |m_2|^2 - |m_1|^2 + 1}{\langle m \rangle + \langle m_2 \rangle + \langle m_1 \rangle}$$

$$\gtrsim \frac{|m| |m_2| (1 - \cos \theta)}{\langle m_1 \rangle}$$

$$\theta = \angle(m_2, -m)$$

Case 1:  $1 - \cos \theta \gtrsim 1$ . (large angle) (60)  
non-resonant

$$I \lesssim \sum_{n=n_1+n_2} \frac{1}{\langle m \rangle^3 \langle m_1 \rangle \langle m_2 \rangle^3} \quad \langle m_i \rangle \sim \max(\langle m \rangle, \langle m_i \rangle)$$


$$\lesssim \langle m \rangle^{-4+}$$

Case 2:  $1 - \cos \theta \ll 1$  (close to collinear) nearly resonant

$$\Rightarrow 0 \leq \theta \ll 1.$$

$$\Rightarrow 1 - \cos \theta \sim \theta^2 \ll 1.$$

Dyadically decompose  $|n_2| \sim N_2$ ,  $N_2 \geq 1$ , dyadic



For fixed  $m \in \mathbb{Z}^3$ ,

$n_2$  is constrained to a cone  $C$

$$|n_2| \sim N_2$$

$$\text{height} \sim N_2 \cos \theta \sim N_2$$

base disc of radius

$$\sim N_2 \sin \theta \sim N_2 \theta.$$

$$\Rightarrow \text{vol}(C) \sim N_2^3 \theta^2$$

$$\Rightarrow I \leq \sum_{\substack{N_2 \geq 1 \\ \text{dyadic}}} \frac{1}{\langle m \rangle^3 \max(\langle m \rangle, N_2) N_2^3 \theta^2} \cdot N_2^3 \theta^2$$

$$\lesssim \langle m \rangle^{-4+}$$



• Prop 15 on  $\gamma = \gamma \oplus 1$

• The proof proceeds analogously to that of Prop 14 but a bit more complicated

New difficult term.

$\gamma > 0$   
small

$$\sum_{\substack{n_2 \in \mathbb{Z}^3 \\ |n+n_2| \wedge |n_2| \\ |n| \ll |n_2|^\gamma}} \frac{\sin(t-t')(\langle n+n_2 \rangle - \langle n_2 \rangle)}{\langle n \rangle^2 \langle n+n_2 \rangle \langle n_2 \rangle^3} \left( \begin{array}{l} \text{WTS} \\ \lesssim \langle n \rangle^{-3} \end{array} \right)$$

This condition allows us to rewrite the sum as

$$\sum_{\substack{n_2 \in \mathbb{Z}^3 \\ |n| \ll |n_2|^\gamma}} \leftarrow \text{we can use symmetrization} \\ n_2 \leftrightarrow -n_2$$

$$\text{Let } \Theta^\pm(n, n_2) = \langle n \pm n_2 \rangle - \langle n_2 \rangle \mp \frac{\langle n, n_2 \rangle}{\langle n_2 \rangle} \\ = \mathcal{O}\left(\frac{\langle n \rangle^2}{\langle n_2 \rangle}\right)$$

$$\text{Sum} = \sum_{\substack{n_2 \in \mathbb{Z}^{3/2} \\ |n| \ll |n_2|^\gamma}} \frac{1}{\langle n \rangle^2 \langle n+n_2 \rangle \langle n_2 \rangle^2} \\ \times \left[ \sin(t-t') \left( \frac{\langle n, n_2 \rangle}{\langle n_2 \rangle} + \Theta^+(n, n_2) \right) \right. \\ \left. - \sin(t-t') \left( \frac{\langle n, n_2 \rangle}{\langle n_2 \rangle} - \Theta^-(n, n_2) \right) \right]$$

$$\text{MVT} \\ \lesssim \sum_{|n| \ll |n_2|^\gamma} \frac{1}{\langle n \rangle^2 \langle n_2 \rangle^3} \left( \frac{\langle n \rangle^2}{\langle n_2 \rangle} \right)^f \text{ for any } f \in [0, 1].$$

$$\lesssim \langle n \rangle^{3+} \quad \square$$



On Prop 18 for  $J \leq \infty$

(63)

The difficult part  $A^{(2)}$

- symmetrization:  $m_2 \leftrightarrow -m_2$
- integration by parts in time  
to handle

$$\sin(t-t')(\langle m+m_2 \rangle + \langle m_2 \rangle)$$